



**Supplemental
Labeling**

EPA Reg. No. 241-379

Expires May 6, 2006

For Postemergence Use in Snap Beans in Delaware, Idaho, Illinois, Indiana, Iowa, Maryland, Michigan, Minnesota, New York, Oregon, Pennsylvania, Virginia, Washington and Wisconsin Only

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING INFORMATION IN THE RAPTOR® HERBICIDE LEAFLET LABEL BEFORE USING. REFER TO THE RAPTOR LEAFLET LABEL FOR FIRST AID AND WORKER PROTECTION STANDARD REQUIREMENTS.

DIRECTIONS FOR USE

READ THE LABEL AFFIXED TO THE CONTAINER FOR RAPTOR HERBICIDE BEFORE APPLYING. USE OF RAPTOR HERBICIDE ACCORDING TO THIS SUPPLEMENTAL LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR RAPTOR HERBICIDE.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of pesticide application.

When **Raptor** is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages may be exceeded. **Raptor** cannot be mixed with any product containing a label prohibiting such mixtures. Do not use **Raptor** other than in accordance with the instructions set forth on this label. Keep containers closed to avoid spills and contamination.

DO NOT apply to snap beans that have been seriously injured from applications of soil-applied herbicides such as Dual® Magnum, Eptam®, or Cobra® herbicides.

THIS PRODUCT WHEN USED ON SNAP BEANS MAY LEAD TO CROP INJURY, LOSS, OR DAMAGE. BASF RECOMMENDS THAT THE USER AND/OR GROWER TEST THIS PRODUCT IN ORDER TO DETERMINE ITS SUITABILITY FOR SUCH INTENDED USE.

GENERAL PRECAUTIONS

DO NOT make more than one application of **Raptor® herbicide** per year.

Allow at least 30 days between application and harvest of snap beans

Only rotational crops harvested at maturity may be used for feed or food.

In the event of a crop loss due to weather, edible legumes, **CLEARFIELD*** canola, **CLEARFIELD*** wheat, or soybeans can be replanted. Do not make an additional application of **Raptor**.

If arid conditions occur during the year of application, rotational crop injury may occur.

Ensure spray drift to non-target species does not occur.

DO NOT apply **Raptor** in any manner not specifically described in this label.

DO NOT apply this product through any type of irrigation system.

When applied by either land or air, **Raptor** spray drift or other indirect contact may injure sensitive crops including non-imidazolinone tolerant wheat or canola, sugarbeets, and leafy vegetables.

Spray equipment used for **Raptor** application must be drained and thoroughly cleaned with water before being used to apply other products.

DISCLAIMER

BASF MAKES THIS PRODUCT AVAILABLE TO THE USER AND/OR GROWER SOLELY TO THE EXTENT THAT THE BENEFIT AND UTILITY, IN THE SOLE OPINION OF THE USER AND/OR GROWER, OUTWEIGH THE EXTENT OF POTENTIAL INJURY ASSOCIATED WITH THE USE OF THIS PRODUCT. THE DECISION TO USE OR NOT TO USE THIS HERBICIDE MUST BE MADE BY EACH INDIVIDUAL **RAPTOR** USER AND/OR GROWER ON THE BASIS OF POSSIBLE CROP INJURY FROM **RAPTOR**, THE SEVERITY OF WEED INFESTATION, THE COST OF ALTERNATIVE WEED CONTROLS, AND OTHER FACTORS. BASF INTENDS THAT BECAUSE OF THE RISK OF FAILURE TO PERFORM OR CROP DAMAGE THAT ALL SUCH USE IS AT THE USER'S AND/OR GROWER'S RISK. BASF DISCLAIMS ANY LIABILITY FOR CLAIMS, CAUSES OF ACTION, FINES, PENALTIES, DAMAGES, INCLUDING CONSEQUENTIAL INCIDENTS AND DAMAGES, LOSSES, LIABILITIES, JUDGEMENTS, AND EXPENSES ARISING OUT OF OR RELATING TO INJURY TO PERSONS, CROPS, OR PROPERTY RESULTING FROM THE USE OF **RAPTOR** ON EDIBLE LEGUMES CONTRARY TO THE LABEL INSTRUCTIONS.

GENERAL INFORMATION

Use Raptor ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Occasionally, internode shortening and/or temporary yellowing of crop plants may occur following **Raptor** applications in snap beans. These effects can be more pronounced if crops are growing under stressful environmental or hot and humid conditions. These effects occur infrequently and are temporary. Normal growth and appearance should resume within days.

APPLICATION INSTRUCTIONS

Early Postemergence Applications: Apply **Raptor** at the broadcast rate of 4 ounces per acre, tank mixed with **Basagran® herbicide** at 6 to 16 ounces per acre. When used in snap beans, **Raptor** must be applied with **Basagran** to minimize crop response. **Basagran** applications at rates higher than 16 ounces per acre may reduce grass control.

Apply to crop and weeds that are actively growing. Apply **Raptor + Basagran** postemergence to snap beans in the first to second trifoliolate leaf stage and weeds that are less than 3 inches tall. Applications before the first trifoliolate leaf stage may result in increased crop response. **DO NOT** apply **Raptor® herbicide + Basagran® herbicide** to snap beans during flowering.

For use in Delaware, Illinois, Indiana, Iowa, Maryland, Michigan, Minnesota, New York, Pennsylvania, Virginia and Wisconsin, a non-ionic surfactant must be added to the spray solution.

The non-ionic surfactant must contain at least 80% active ingredient and should be used at a rate of 1 quart per 100 gallons of spray solution.

For use in Idaho, Oregon and Washington, a non-ionic surfactant and nitrogen fertilizer must be added to the spray solution. The non-ionic surfactant must contain at least 80% active ingredient and should be used at a rate of 1 quart per 100 gallons of spray solution. Alternatively, a crop oil concentrate (COC) or methylated seed oil (MSO) can be used. Use COC at a rate of 1 gallon per 100 gallons of spray solution. Use MSO at a rate of 1 - 2 gallons per 100 gallons of spray solution. Recommended nitrogen based fertilizers include liquid fertilizers such as 28-0-0, 32-0-0, or 10-34-0 at the rate of 2.5 gallons per 100 gallons of spray solution. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 12 – 15 pounds per 100 gallons of spray solution.

Raptor tank mixes with any pesticide other than **Basagran** are not recommended. Certain insecticide and herbicide tank mixes with **Raptor** in snap beans have shown unacceptable crop response.

Raptor may be applied a minimum of one hour before rainfall or overhead irrigation.

BROADLEAF WEEDS CONTROLLED BY RAPTOR

	<u>Application Rate</u>
	Raptor at 4 oz./A + Basagran at 6 – 16 oz./A
	Maximum Weed Size (inches)
Bedstraw	3
Beet, wild	3
Buttercup	3
Chickweed, common	3
Jimsonweed	3
Mustard,	
tumble	3
wild	3
black	3
Nightshade,	
black	3
Eastern black	3
Hairy	3
Pennycress, field	3
Pigweed,	
Redroot	3
Smooth	3
Spiny	3
Puncturevine	3
Radish, wild	3
Shepherdspurse	3
Tansymustard, green	3

**BROADLEAF WEEDS SUPPRESSED BY
RAPTOR® HERBICIDE**

<u>Application Rate</u>	
Raptor at 4 oz./A + Basagran at 6 – 16 oz./A	
Maximum Weed Size (inches)	
Buckwheat, wild	3
Cocklebur, common	3
Lambsquarters, common	3
Knotweed, prostrate	3
Kochia*	3
Lettuce, miners	3
Morningglory	
Entireleaf	3
Ivyleaf	3
Smallflower	3
Tall	3
Purslane, common	3
Rocket, London	3
Smartweed	
Ladysthumb	3
Pennsylvania	3
Spurge, prostrate	3

***Raptor** control non-ALS resistant kochia only.

GRASS WEEDS CONTROLLED BY RAPTOR® herbicide

<u>Application Rate</u>	
Raptor at 4 oz./A + Basagran at 6 – 16 oz./A	
Maximum Weed Size (inches)	
Barnyardgrass	3
Blackgrass	3
Brome,	
downy	3
cheat	3
Japanese	3
Canarygrass, littleseed	3
Cereals, volunteer	
Barley	3
Oat	3
Wheat (non- CLEARFIELD*)	3
Darnel, Persian	3
Foxtail,	
Giant	3
Green	3
Yellow	3
Jointed goatgrass	3
Oats, wild	3
Ryegrass, Italian	3
Shattercane	3
Volunteer corn*	2-8

* Except imidazolinone tolerant corn

GRASS WEEDS SUPPRESSED BY RAPTOR

<u>Application Rate</u>	
Raptor at 4 oz./A + Basagran at 6 – 16 oz./A	
Maximum Weed Size (inches)	
Johnsongrass, rhizome	3
Crabgrass,	
Large	3
Smooth	3
Sedges	
Purple	3
Yellow	3
Quackgrass	3

ROTATIONAL CROP RESTRICTIONS

See main **Raptor** label for rotational plant-back intervals following applications of **Raptor**.

©2004 BASF Corporation
All rights reserved

Basagran, CLEARFIELD and Raptor are registered trademarks of BASF.
Dual Magnum and Eptam are registered trademarks of Syngenta.
Cobra is a registered trademark of Valent USA Corp.

000241-00379.20041029b.**NVA 2004-04-133-0284**
Supersedes NVA 2004-04-133-0093

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709



The Chemical Company