

Endura™

fungicide

Supplemental Labeling

For use in head and stem brassicas and leafy brassica greens.

EPA Reg. No. 7969-197

Active Ingredients:

Boscalid - (3-pyridinecarboxamide, 2-chloro-N-(4'-chloro(1,1'-biphenyl)-2-yl)) 70.0%

Inert ingredients 30.0%

Total 100.0%

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Refer to the **Endura™ fungicide** main label for precautionary statements, first aid and personal protective equipment requirements. This supplemental label must be in the user's possession during application.

General Information

Endura provides optimum disease control when applied in a regularly scheduled protective fungicide program and is used in a spray program that rotates fungicides with different modes of action. Refer to the **Endura** main label for general resistance management information and to the crop specific use recommendations and restrictions found in this label.

Application Information

Apply **Endura** according to the rate, timing, resistance management and adjuvant use recommendations in the **Crop Specific Use Directions** (Table A) in this label. Observe the additional general recommendations on application instructions, additive use and mixing order on the **Endura** main label.

Endura may be applied by ground sprayer, aerial equipment or through sprinkler irrigation systems. Refer to the **Endura** main label for specific instructions on these methods.

Restrictions and Limitations

Endura is not for use in greenhouse or transplant production systems.

Follow the restrictions and limitations outlined in the **Crop Specific Restrictions and Limitations** table (Table B) in this label for:

- Minimum pre-harvest interval (PHI)
- Maximum rate per acre
- Maximum number of applications
- Maximum rate per season
- Livestock grazing or feeding restrictions
- Aerial application restrictions

BASF

Table A– **Endura**™ fungicide Crop-Specific Use Directions

Brassica Leafy Vegetable Crop	Target Diseases	Use Rate per Application	Maximum Number of Applications per Season	Maximum Rate per Season	Minimum Time from Application to Harvest (PHI)
Head and Stem Brassicas: Broccoli Chinese broccoli (gai lon) Brussels sprouts Cabbage Chinese cabbage (napa) Chinese mustard (gai choy) Cauliflower Cavalo broccolo Kohlrabi	Alternaria blight (Black spot) (<i>Alternaria</i> spp.) Grey mold (<i>Botrytis cinerea</i>) Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i> , <i>S. minor</i>) Suppression only: Powdery mildew (<i>Erysiphe polygoni</i>) Rhizoctonia bottom rot (<i>Rhizoctonia solani</i>)	6 to 9 oz. per acre	2	18 oz. per acre	0 days
<p>Application Directions: Begin applications of Endura prior to disease development and continue on a 7 to 14 day interval. Use the higher rate and the shorter interval when disease pressure is high.</p> <p>Resistance Management: To limit the potential for development of resistance, do not make more than two (2) applications of Endura per season.</p>					
Leafy Brassica Greens: Broccoli raab (rapini) Chinese cabbage (bok choy) Collards Kale Mizuna Mustard greens Mustard spinach Rape greens	Alternaria blight (Black spot) (<i>Alternaria</i> spp.) Grey mold (<i>Botrytis cinerea</i>) Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i> , <i>S. minor</i>) Suppression only: Powdery mildew (<i>Erysiphe polygoni</i>) Rhizoctonia bottom rot (<i>Rhizoctonia solani</i>)	6 to 9 oz. per acre	2	18 oz. per acre	14 days
<p>Application Directions: Begin applications of Endura prior to disease development and continue on a 7 to 14 day interval. Use the higher rate and the shorter interval when disease pressure is high.</p> <p>Resistance Management: To limit the potential for development of resistance, do not make more than two (2) applications of Endura per season.</p>					

Table B – **Endura™** fungicide Restrictions and Limitations

Crop	Minimum Time from Application to Harvest (PHI) (days)	Maximum Rate per Acre per Application (oz.)	Maximum Number of Applications per Season	Maximum Rate per Acre per Season (oz.)	Aircraft Application
Head and Stem Brassicas ¹ : Cabbage Broccoli Cauliflower	0	9	2	18	Yes
Leafy Brassica Greens ¹	14	9	2	18	Yes
¹ For a complete list of crops within a crop group, see Table A. Endura™ fungicide Crop-Specific Use Directions.					

Endura is a trademark of **BASF**

© 2004 BASF Corporation
All Rights Reserved

007969-00197.20040429.NVA 2004-04-201-0085.pdf
Supersedes: NVA 2003-04-201-0029

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709

BASF